

Quantal corrections to mean-field dynamics including pairing

Denis Lacroix* and Danilo Gambacurta

GANIL, CEA and IN2P3, Boîte Postale 5027, 14076 Caen Cedex, France

Sakir Ayik

Physics Department, Tennessee Technological University, Cookeville, TN 38505, USA

Extending the stochastic mean-field model by including pairing, an approach is proposed for describing evolutions of complex many-body systems in terms of an ensemble of Time-Dependent Hartree-Fock Bogoliubov trajectories which is determined by incorporating fluctuations in the initial state. Non-linear evolution of the initial fluctuations provides an approximate description of quantal correlations and fluctuations of collective observables. Since the initial-state fluctuations break the particle-number symmetry, the dynamical description in which pairing correlations play a crucial role is greatly improved as compare to the mean-field evolution. The approach is illustrated for a system of particles governed by a pairing Hamiltonian.

PACS numbers: 24.10.Cn, 05.40.-a, 05.30.Rt

Keywords: Pairing, Many-body dynamics, symmetry breaking, stochastic methods

Under certain conditions, it is possible to provide an approximate description for quantal evolution of a system in terms of an ensemble of classical trajectories with proper choice of initial conditions. This aspect appears naturally in the path integral formulation of quantum dynamics and has been recognized in refs. [1, 2]. In recent years, this idea has been pushed forward to improved the mean-field description of many-body interacting systems [3, 4]. Mean-field description cannot describe essential quantal effects associated to collective motion and severely underestimates fluctuations of collective observables. By considering an ensemble of mean-field trajectories with a specific choice of fluctuations (quantal zero-point and thermal) in the initial state, it is possible to overcome some of these shortcomings. Several applications, especially in the nuclear physics context [5], have shown that this approach can improve the mean-field description by including important dissipative aspects in transport properties in heavy-ion collisions. More recently, we illustrated that such an approach, in addition to fluctuations, can accurately tackle the problem of symmetry breaking close to bifurcation point in collective energy landscape [6]. Up to now, the stochastic mean-field (SMF) approach with initial fluctuations has been developed starting from a Time-Dependent Hartree-Fock (TDHF) version of the mean-field. Nowadays, there are increasing interests in the treatment of pairing to describe evolution of strongly interacting Fermi liquids [7–11] employing the Time-Dependent Hartree-Fock Bogoliubov (TDHFB) approach or its simplified BCS limit. While the TDHFB theory provides an important improvement beyond TDHF, it still suffers from the above quoted limitations: i.e. underestimation of quantum collective fluctuations and impossibility to spontaneously break symmetries. Due to successful description of the SMF approach in geometric symmetry breaking, it is rather tempting to introduce quantum fluctuations through ini-

tial sampling to treat pairing where the $U(1)$ symmetry breaking plays an important role.

In this work, we present an extension of the SMF approach by incorporating pairing into the description. In the standard TDHFB theory, evolution of the generalized density matrix $\mathcal{R}(t)$ is given by

$$i\hbar \frac{d}{dt} \mathcal{R}(t) = [\mathcal{H}(\mathcal{R}), \mathcal{R}(t)]. \quad (1)$$

$\mathcal{R}(t)$ contains both normal $\rho_{ij} = \langle a_j^\dagger a_i \rangle$ and anomalous $\kappa_{ij} = \langle a_j a_i \rangle$ density matrix components. Here, $(a_i^\dagger, a_i)$ are creation and annihilation operators of a given single-particle basis. The quantity $\mathcal{H}(\mathcal{R})$ is the generalized mean-field Hamiltonian containing both the mean-field $h[\rho]$ and the pairing field $\Delta[\kappa]$ [12]. Eq. (1) is generally solved either starting from a density associated to a quasi-particle vacuum or a statistical ensemble of quasi-particles. It is then convenient to introduce associated quasi-particle creation operators, defined through $\gamma_\alpha^\dagger = \sum_i U_{i\alpha}^* a_i + V_{i\alpha}^* a_i^\dagger$. In quasi-particle representation, normal density matrix is diagonal and expectation value of anomalous density matrix vanishes,

$$\rho_{\alpha\beta} = \langle \gamma_\beta \gamma_\alpha^\dagger \rangle = \delta_{\alpha\beta} f_\alpha; \quad \kappa_{\alpha\beta} = \langle \gamma_\beta \gamma_\alpha \rangle = 0, \quad (2)$$

where f_α denotes the occupation numbers of quasi-particle states [12, 13] and the expectation values are taken with respect to the quasi-particle vacuum. As a result, $\mathcal{R}(t)$ takes diagonal form

$$\mathcal{R}(t) = \sum_\alpha |W_\alpha(t)\rangle f_\alpha \langle W_\alpha(t)| + |V_\alpha(t)\rangle (1 - f_\alpha) \langle V_\alpha(t)| \quad (3)$$

¹ In the following, Greek index will be reserved to quasi-particles while latin index will refers to particles.

where vectors $|V_\alpha(t)\rangle$ and $|W_\alpha(t)\rangle$ are the eigenvectors of $\mathcal{R}(t)$ whose expressions can be found in [12]. In following, for notation compactness, we will use $|S_\alpha(t)\rangle$ for both types of states. Then, in the TDHFB dynamics, we need to evolve state vectors according to $i\hbar\partial_t|S_\alpha\rangle = \mathcal{H}(\mathcal{R})|S_\alpha\rangle$, while keeping the quasi-particle occupation fixed. In the stochastic extension of TDHFB, quantal zero-point fluctuations (and thermal fluctuations) in the initial state are incorporated into the description in a similar manner to the SMF approach developed in [3]. Initial fluctuations are simulated by generating an ensemble of initial density matrices. Each member of the initial density matrix, labelled by (n) is written as,

$$\mathcal{R}^{(n)}(t_0) = \sum_{\alpha\beta} |S_\alpha(t_0)\rangle \mathcal{R}_{\alpha\beta}^{(n)} \langle S_\beta(t_0)|. \quad (4)$$

where summations over α and β run over a complete set of state vectors. The statistical properties of the elements of density matrix $\mathcal{R}_{\alpha\beta}^{(n)}(t_0)$ is specified in terms of statistical properties of the normal $\rho_{\alpha\beta}^{(n)}(t_0)$ and anomalous $\kappa_{\alpha\beta}^{(n)}(t_0)$ density matrices. Elements of the normal and the anomalous density matrices are uncorrelated Gaussian random numbers with the mean values,

$$\overline{\rho_{\alpha\beta}^{(n)}} = \delta_{\alpha\beta} f_\alpha, \quad \overline{\kappa_{\alpha\beta}^{(n)}} = 0. \quad (5)$$

and the second moments defined by,

$$\overline{\delta\rho_{\alpha\beta}^{(n)} \delta\rho_{\alpha'\beta'}^{(n)*}} = \frac{1}{2} \delta_{\alpha\alpha'} \delta_{\beta\beta'} [f_\alpha(1-f_\beta) + f_\beta(1-f_\alpha)], \quad (6)$$

$$\overline{\delta\kappa_{\alpha\beta}^{(n)} \delta\kappa_{\alpha'\beta'}^{(n)*}} = \frac{1}{2} \delta_{\alpha\alpha'} \delta_{\beta\beta'} [f_\alpha f_\beta + (1-f_\alpha)(1-f_\beta)]. \quad (7)$$

In these expressions, $\delta\rho^{(n)}$ and $\delta\kappa^{(n)}$ are the fluctuating parts of the density matrices,

$$\rho^{(n)} = \overline{\rho^{(n)}} + \delta\rho^{(n)}, \quad \kappa^{(n)} = \overline{\kappa^{(n)}} + \delta\kappa^{(n)} \quad (8)$$

and notation $\overline{X}$ indicates the average over the initial ensemble. We note that in the SMF approach without pairing only fluctuations in the particle-hole channels are included. In the extension including pairing, additional fluctuations originating from particle-particle and hole-hole channels appears, Eq. (7). This is an important new aspect to explore initial conditions with non-zero anomalous densities.

In the novel stochastic approach, an event of the ensemble of generalized density matrices is expressed as,

$$\mathcal{R}^{(n)}(t) = \sum_{\alpha\beta} |S_\alpha^{(n)}(t)\rangle \mathcal{R}_{\alpha\beta}^{(n)}(t_0) \langle S_\beta^{(n)}(t)|, \quad (9)$$

where states in an event evolve with its own self-consistent generalized Hamiltonian according to the TDHFB equation of motion, $i\hbar\partial_t|S_\alpha^{(n)}\rangle = \mathcal{H}(\mathcal{R}^{(n)})|S_\alpha^{(n)}\rangle$. In

the TDHFB framework, we can consider generalized operators that can create or annihilate two particles,

$$\hat{Q} = \sum_{ij} Q_{ij}^{11} a_i^\dagger a_j + \sum_{ij} \left(Q_{ij}^{20} a_j a_i + Q_{ji}^{20*} a_i^\dagger a_j^\dagger \right). \quad (10)$$

The expectation value of such an operator in each event generated in the stochastic TDHFB approach reads

$$Q^{(n)}(t) = \text{Tr}(Q^{11}\rho^{(n)}(t)) + 2\Re\text{Tr}(Q^{20}\kappa^{(n)}(t)). \quad (11)$$

Then, we determine the average evolution $\overline{Q}(t)$ by taking the average over the ensemble generated in the simulations. Since stochastic approach incorporates correlations beyond the mean-field description, the ensemble average value $\overline{Q}(t)$, in general, is different than the expectation value of the operator in the standard TDHFB approach. Dispersion of the observable $\hat{Q}$ is calculated using the classical formula,

$$\sigma_Q^2(t) = \overline{(Q^{(n)}(t) - \overline{Q}(t))^2}. \quad (12)$$

We note that, fluctuations (6-7) are chosen to insure the initial ensemble average of the first and second moments equal to those obtained from the quasi-particle density matrix at zero or finite temperature.

In order to illustrate the powerfulness of the approach, we consider a many-body system governed by a pairing Hamiltonian. In the model, there are K single-particle levels, labelled by i . Each level is associated with the energy ε_i and has a degeneracy $2\Omega_i$. It is assumed that each time a state i is present, its time-reversed states $\bar{i}$ is also present with same energy. We define number operators for each energy level as $\hat{N}_i = \sum_p^{\Omega_i} (a_{p,i}^\dagger a_{p,i} + a_{\bar{p},i}^\dagger a_{\bar{p},i})$. In addition, pair creation/annihilation operators $\hat{S}_i^+ = \sum_p^{\Omega_i} a_{p,i}^\dagger a_{\bar{p},i}^\dagger$, $\hat{S}_i^- = (\hat{S}_i^+)^\dagger$ are introduced. The pairing Hamiltonian of the system is given by [14]

$$H = \sum_{i=1}^K \varepsilon_i \hat{N}_i + \sum_{ij} G_{ij} \hat{S}_i^+ \hat{S}_j^-. \quad (13)$$

From the set of operators, one can construct three quasi-spin components $\hat{S}_i^x = (\hat{S}_i^+ + \hat{S}_i^-)/2$, $\hat{S}_i^y = (\hat{S}_i^+ - \hat{S}_i^-)/2i$ and $\hat{S}_i^z = (\hat{N}_i - \Omega_i)/2$, which form a standard SU(2) algebra. The eigenstates of H can be determined by direct diagonalization [15] giving access to the exact static and dynamical properties.

Within TDHFB, evolutions of the expectation values ($S_i^x(t)$, $S_i^y(t)$, $S_i^z(t)$) of the quasi-spin components are determined by a set of coupled equations ($\hbar = 1$),

$$\frac{d}{dt} \begin{pmatrix} S_i^x(t) \\ S_i^y(t) \\ S_i^z(t) \end{pmatrix} = \begin{pmatrix} 0 & -2\varepsilon_i & +2\Delta_i^y \\ 2\varepsilon_i & 0 & -2\Delta_i^x \\ -2\Delta_i^y & 2\Delta_i^x & 0 \end{pmatrix} \begin{pmatrix} S_i^x(t) \\ S_i^y(t) \\ S_i^z(t) \end{pmatrix} \quad (14)$$

where $\Delta_i^{x,y}(t) = \sum_k G_{ik} S_i^{x,y}(t)$. These equations of motion are quite general and except that single-particle energy are fixed in time, such equations can be found in

Fermi gas dynamics simulations on lattice with contact interaction [7] or in nuclear physics context [10, 11].

In order to illustrate some limitations of TDHFB, let us suppose that the initial state of the system is described by a Slater determinant of single-particle wave functions. Since the Hamiltonian (13) contains two-body interactions, this state is not an eigenstate of the pairing Hamiltonian and it is expected to evolve in time. However, it turns out that this state is a stationary solution of the TDHFB equation. Indeed, since the state is an eigenstate of the particle number, the expectation values of quasi-spins vanish $S_i^x(t_0) = S_i^y(t_0) = 0$. Hence, we deduce that all quasi-spin components remain constant in time. Therefore, the TDHFB approach is unable to describe correlations that built up in time and leads to the departure from the independent particle approach. This failure stems from the fact that a symmetry that exists initially, here the $U(1)$ symmetry, cannot be spontaneously broken in the mean-field approach. In the stochastic TDHFB approach, the situation is different. Expectation values of quasi-spin components are still determined by the same set of equations (14). However, the initial condition is not determined in a deterministic manner, but is specified in terms of distributions of $(S_i^{x(n)}, S_i^{y(n)}, S_i^{z(n)})_{i=1,K}$. Following from the basic postulate of the approach, (6) and (7), the initial distributions of quasi-spin components are uncorrelated random Gaussian numbers. Their mean values are determined by $S_i^{x(n)} = S_i^{y(n)} = 0$ while $S_i^{z(n)} = +1/2$ or $-1/2$ for occupied and unoccupied levels respectively. Identifying the quasi-particle vacuum with a Slater determinant, variances of quasi-spin distributions are given by,

$$\overline{S_i^{z(n)} S_j^{z(n)}} = \overline{S_i^{z(n)} S_j^{x(n)}} = \overline{S_i^{z(n)} S_j^{y(n)}} = \overline{S_i^{x(n)} S_j^{y(n)}} = 0$$

and

$$\overline{S_i^{x(n)} S_j^{x(n)}} = \overline{S_i^{y(n)} S_j^{y(n)}} = \frac{1}{4} \delta_{ij} \Omega_i.$$

Consequently, for each level i , the z component of quasi-spin is initially a non-fluctuating quantity, while for each i , x and y components $(S_i^{x(n)}, S_i^{y(n)})$ are specified by uncorrelated real Gaussian distributions. Solution of Eq. (14) is determined by generating an ensemble of events according to the initial distributions of quasi-spin components. Solution is not a deterministic final state, but consists of a distribution of final states. Note that the mean number of particle is conserved event by event while the total energy is conserved in average.

In the illustrations below, we consider $K = 10$ doubly-degenerated levels ($\Omega_i = 1$) with constant level spacing $\Delta\epsilon$ between adjacent levels, and constant pairing interaction $G_{ik} = G$. A system of $N = 10$ particles occupy initially the five lowest energy levels. The exact evolution is obtained by decomposing the initial state in terms of eigenstates of the pairing Hamiltonian. The results

are compared with stochastic simulations by generating $N_{\text{evt}} = 2 \cdot 10^5$ number of events. We obtain solutions of Eq. (14) by employing a Runge-Kutta 2 algorithm with a numerical time step $\Delta t = 0.005/\Delta\epsilon$. In Fig. 1 the quantity $D(t)$, which illustrates the time-scale associated to the departure from the independent particle,

$$D(t) = \sum_i \Omega_i n_i(t) (1 - n_i(t)) \quad (15)$$

is plotted as a function of time for different coupling strength $G/\Delta\epsilon$. Here $n_i(t)$ are the single-particle occupation numbers. In this figure, symbols show the exact results and solid lines denote the simulations of stochastic TDHFB approach. In the stochastic TDHFB, the average occupation numbers are determined by the following ensemble averages $n_i(t) = 1/2 + \overline{S_i^{z(n)}}/\Omega_i$.




FIG. 1: (color online) Exact evolution of $D(t)$ for $G/\Delta\epsilon = 0.05$ (black triangles), 0.5 (blue squares) and 1 (red circles). The results obtained by averaging over TDHFB trajectories are shown respectively by black long dashed line, blue short dashed line and red solid line. Note that curves corresponding to $G/\Delta\epsilon = 0.05$ have been multiplied by 50 . The time is presented in $(\Delta\epsilon)^{-1}$ unit.

The stochastic approach with pairing is able to incorporate non-trivial dissipative effects due to the coupling between complex internal degrees of freedom. Incorporating non-trivial initial fluctuations in particle-particle (pp) and hole-hole (hh) channels is able in all cases to describe short time dynamics very well. When the interaction strength increases, evolution of $D(t)$ becomes more complex and tends to oscillate around an average value. The stochastic approach is accurate not only for short time behavior, but also gives a quite reasonable description of the asymptotic behavior although the evolutions appear more damped than the exact solutions. In the weak coupling regime $G/\Delta\epsilon = 0.05$, the exact dynamics exhibits a periodic evolution with a rather short Poincaré recurrence time that is perfectly reproduced by the average TDHFB evolution. We note that, with the present parameter value for the Hamiltonian, the BCS

threshold is at $G_{\text{cr}}/\Delta\varepsilon \simeq 0.3$, i.e. above this coupling strength, a $U(1)$ symmetry breaking mean-field solutions exists. It is important to recall that the standard TDHFB gives $D(t) = 0$. Similarly, the previously proposed version of SMF approach that was not exploring the possibility to break the particle number symmetry also leads to $D(t) = 0$.

As another illustration, the two-body quantity $\sigma_{\pm}(t) = \sum_{ij} \langle \hat{S}_i^+ \hat{S}_j^- \rangle(t)$ is considered. Fig. 2 shows this quantity as a function of time for the exact and the stochastic TDHFB approach for different values of coupling strength $G/\Delta\varepsilon$. Symbols indicate the exact evolutions and the solid lines are the results of the stochastic TDHFB simulations from the expression $\sigma_{\pm}(t) = \sum_{ij} \left(\overline{S_i^{x(n)}(t) S_j^{x(n)}(t)} + \overline{S_i^{y(n)}(t) S_j^{y(n)}(t)} \right)$. Here also, we




FIG. 2: (color online) Comparison of the exact and average evolution of $\sigma_{\pm}(t)$ for different coupling strengths. Conventions are the same as in Fig. 1. Note that curves corresponding to $G/\Delta\varepsilon = 0.05$ have been multiplied by a factor 5.

observe the stochastic approach reproduces perfectly the exact evolution of quantal fluctuations during relatively short time scale for all coupling strengths, and provides a reasonable description of the gross properties (i.e. time average behavior) of the fluctuations over a long time scale for all coupling strengths

In the present work, we propose a stochastic extension of the mean-field dynamics including pairing correlations. In this approach, quantal zero-point (and possibly thermal) fluctuations in the initial state are incorporated by simulating an ensemble of events by constructing a suitable distribution of the initial states. Observable quantities are determined by averaging over the generated ensemble. The previous illustrations of the approach without pairing [3, 6] and the present application including the pairing correlations show that the stochastic approach reproduces a reasonable description of the

gross properties exact quantal dynamics. These illustrations indicate a strong support for the fact the stochastic extension of the mean-field without and with pairing provide a powerful tool for an approximate description of the correlations and quantal fluctuations of collective motion in normal and superfluid mesoscopic systems, beyond the mean-field description. The TDHFB approach has recently become a widely employed tool to describe Fermi liquids appearing in nuclear, condensed matter or atomic physics. With rapid progress of computational powers, it is not anymore unreasonable to simulate thousands of independent TDHFB events, and consequently the stochastic approach proposed in this work may provided as the next step to further progress in the description of transport properties of such physical systems.

Acknowledgment

S. A. gratefully acknowledge GANIL and IPN Orsay for hospitality extended to him during his visits. This work is supported in part by the US DOE Grant No. DE-FG05-89ER40530.

* Electronic address: lacroix@ganil.fr

- [1] M. F. Herman and E. Kluk, Chem. Phys. **91**, 27 (1984).
- [2] K. G. Kay, J. Chem. Phys. **100**, 4432 (1994); **101**, 2250 (1994).
- [3] S. Ayik, Phys. Lett. **B658**, 174 (2008).
- [4] D. Lacroix, S. Ayik, and Ph. Chomaz, Prog. Part. Nucl. Phys. **52**, 497 (2004).
- [5] S. Ayik, K. Washiyama, and D. Lacroix, Phys. Rev. **C79**, 054606 (2009). K. Washiyama, S. Ayik, and D. Lacroix, Phys. Rev. **C80**, 031602(R) (2009). S. Ayik, B. Yilmaz and D. Lacroix, Phys. Rev. **C81**, 034605 (2010). B. Yilmaz, S. Ayik, D. Lacroix, and K. Washiyama, Phys. Rev. **C83**, 064615 (2011).
- [6] D. Lacroix, S. Ayik and B. Yilmaz, Phys. Rev. **C 85**, 041602(R) (2012).
- [7] G. Tonini, F. Werner and Y. Castin, Eur. Phys. J. D **39**, 283 (2006).
- [8] B. Avez, C. Simenel, and Ph. Chomaz, Phys. Rev. **C 78**, 044318 (2008).
- [9] I. Stetcu, A. Bulgac, P. Magierski, and K. J. Roche Phys. Rev. **C 84**, 051309, (2011).
- [10] S. Ebata, T. Nakatsukasa, T. Inakura, K. Yoshida, Y. Hashimoto, and K. Yabana, Phys. Rev. **C 82**, 034306 (2010).
- [11] G. Scamps and D. Lacroix, Phys. Rev. **C87**, 014605 (2013).
- [12] J. P. Blaizot and G. Ripka, *Quantum Theory of Finite Systems*, (MIT Press, Cambridge, Massachusetts, 1986).
- [13] A.L. Goodman, Nucl. Phys. **A352**, 30 (1981), A.L. Goodman, Phys. Rev. **C29**, 1887 (1984).
- [14] R. W. Richardson and N. Sherman, Nucl. Phys. **52**, 221 (1964); R.W. Richardson, Phys. Rev. **141**, 949 (1966); J.Math. Phys. **9**, 1327 (1968).

- [15] A. Volya, B. A. Brown, and Z. Zelevinsky, Phys. Lett. **B 509**, 37 (2001); T. Sumaryada and Alexander Volya, Phys. Rev. **C 76**, 024319 (2007).